

NUTRIENTS PER 100 CALORIES (5 FL OZ, PREPARED AS DIRECTED)

PROTEIN	2.45 g	WATER		133 g
FAT	5.46 g	LINOIC ACID		1000 mg
CARBOHYDRATE	10.4 g			

VITAMINS

VITAMIN A	300 IU	NIACIN	1350 mcg
VITAMIN D	60 IU	FOLIC ACID (FOLACIN)	15 mcg
VITAMIN E	1.5 IU	PANTOTHENIC ACID	750 mcg
VITAMIN K	11 mcg	Biotin	4.5 mcg
THIAMIN (VIT. B ₁)	60 mcg	VITAMIN C	
RIBOFLAVIN (VIT. B ₂)	90 mcg	(ASCORBIC ACID)	9 mg
VITAMIN B ₆	60 mcg	CHOLINE	12 mg
VITAMIN B ₁₂	0.45 mcg	inositol	5 mg

MINERALS

CALCIUM	105 mg	COPPER	75 mcg
PHOSPHORUS	75 mg	IODINE	15 mcg
MAGNESIUM	7.5 mg	SELENIUM	1.8 mcg
IRON	1.8 mg	SODIUM	44 mg
ZINC	0.75 mg	POTASSIUM	108 mg
MANGANESE	25 mcg	CHLORIDE	62 mg

INGREDIENTS: 39.2% CORN SYRUP SOLIDS, 14.7% SOY PROTEIN ISOLATE, 11.5% HIGH OLEIC SAFFLOWER OIL, 9.8% SUGAR (SUCCROSE), 8.4% SOY OIL, 7.7% COCONUT OIL, LESS THAN 2% OF: C. COHNII OIL*, M. ALPINA OIL*, BETA-CAROTENE, LUTEIN, LYCOPENE, FRUCTOOLIGOSACCHARIDES, CALCIUM PHOSPHATE, POTASSIUM CITRATE, POTASSIUM CHLORIDE, MAGNESIUM CHLORIDE, SODIUM CHLORIDE, ASCORBIC ACID, CHOLINE CHLORIDE, L-METHIONINE, TAURINE, ASCORBYL PALMITATE, FERROUS SULFATE, m-INOSITOL, MIXED TOCOPHEROLS, ZINC SULFATE, d-ALPHA-TOCOPHERYL ACETATE, L-CARNITINE, NIACINAMIDE, CALCIUM PANTOTHENATE, COPRIC SULFATE, THIAMINE CHLORIDE HYDROCHLORIDE, VITAMIN A PALMITATE, RIBOFLAVIN, PYRIDOXINE HYDROCHLORIDE, FOLIC ACID, POTASSIUM IODIDE, POTASSIUM HYDROXIDE, PHYLOQUINONE, BIOTIN, SODIUM SELEVATE, VITAMIN D₃, AND CYANOCOBALAMIN.

CONTAINS SOY INGREDIENTS.

*Contains no dairy ingredients. Manufactured on dairy equipment.
†SOURCE OF DOCSAHEXAENOIC ACID (DHA) ‡SOURCE OF ARACHIDONIC ACID (ARA)

Abbott Nutrition
Abbott Laboratories
Columbus, Ohio 43219-3034 USA

U.S. Patents 5,221,545; 6,136,958;
6,596,767 and 7,090,879
© 2010 Abbott Laboratories
#55963 38148 8768-01

Similac® formulas
for every baby's nutritional needs

Advance
Complete nutrition for the 1st year
Advance®
Organic

Sensitive
For babies with fussiness, gas, & spit-up
Sensitive
Soy
Spit-Up

Expert Care
For babies needing extra TLC
Alimentum®
NeoSure®
Diarrhea

Plus Similac Go & Grow® for toddlers learning to eat table food

Our Feeding Expert hotline is available to help you with any feeding issues: 800-986-8800

USE AS DIRECTED BY A DOCTOR
Directions for Preparation and Use

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) nipple and ring before use.

Storage

Once mixed, store bottles in refrigerator and feed to baby within 24 hours. Close opened can and store in a cool, dry area (not in refrigerator). Use opened can contents within 1 month.

Warning

Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Never use a microwave to warm formula. Serious burns can result.

Mixing Guide

Measure water	+ Add scoop(s) of unpacked level powder using enclosed scoop	= Finished bottle (approx.)
2 fl oz	1 scoop (8.8 g)	2 fl oz
4 fl oz	2 scoops	4 fl oz
8 fl oz	4 scoops	8 fl oz

1 Can = 15 Bottles
When mixed as directed, makes approx. 90 fl oz of formula.

Have product-related questions?
Call 1-800-515-7677, 8:30 am–5:00 pm,
Eastern time, weekdays.
www.Similac.com

Abbott

WITH IRON | POWDER
BIRTH TO 12 MONTHS
DHA/ARA

Similac

SENSITIVE®

Isomil Soy™

For Fussiness & Gas

#1 Formula for Complete Soy Nutrition

NET WT. 12.4 OZ (352 g)

13-1/16"
3-1/2-1/16"
100mm LINE

Version #11 09/22/08